Review Article:

Viral Encephalitis

DR. B. P. NEPAL*

The organism which slaughtered thousands in India and Nepal last year is Japanese B Encephalitis virus. It is a group B type of arbo virus. It is a small RNA virus and there are many antigenically related types. The insect responsible for the transmission of this virus is *Culex gelidus* in this part of the world. The infection in man is only the spill over of infection in pigs and birds. The disease is infrequent manifestation of human infection. The infection rate in the epidemic area in the children is about 10% but only one in three to five hundred infection results in disease, but in adults one in twentyfive infection results in disease. The epidemic is dreaded to be larger this year as it is larger in alternate years. The epidemic in man usually occurs in August,–September, about one month after it appears in pigs and birds. But it can be less seasonal.

PATHOGENESIS :

When an infected mosquito bites a man, the virus is deposited in the skin where it multiplies. It is then taken by the lymphatics to the lymph nodes, spleen and small blood vessels. It undergoes further multiplication in those organs and then released in blood. Viremia and dissemination to other organs then results. The incubation period is about 10 days. The extrinsic incubation period (in the mosquito) is about one week. The infectivity in the mosquito is retained throughout its life.

PATHOLOGY :

The organ severely damaged is brain. The brain becomes grossly oedematous with occasional small haemorrhages. On microscopic examination, oedema and hyperemia in the lepto-meninges and cortex are seen. Small haemorrhages and arteriolar thrombi are also seen. Perivascular cuffing is characteristic. Evidence of focal neuronal lesions are

* M. O., Palpa Health Club.

also found. There is degeneration of cytoplasm and nucleus. Focal nodules are found with microglial and polymorphonuclear accumulation. The damage is either due to itself or due to immunologically related injury.

In the blood vessels, damage to the vascular endothelial cells are seen. No virus is isolated from the blood vessels. There may be myocarditis and focal hepatic necrosis.

CLINICAL MANIFESTATION :

The symptoms are due to:

- (1) Permanent Neuronal damage
- and (2) Temporary central dysfunction due to anoxia and oedema.

The onset is usually mild. The early symptoms are drowsiness, headache, nausea, vomiting and myalgia. This is called the prodromal stage. This is followed by persistent high fever which cannot be brought down by conventional antipyretics. The temperature is due to anoxia of the thermoregulatory centre. The fever peaks four to five days after the onset and slowly subsides. There is relative bradycardia. The pattern of disease in children is slightly different from that in the adult. Convulsion, disorientation, abnormal reflexes and paralysis of face and extremities are the features in children. In adult, bilateral paresis without sensory changes is the feature. But there may be sensorial changes, spastic rigidity, cerebellar signs and coma which may result in death.

INVESTIGATIONS :

There are no sophisticated clinical investigations to fix the diagnosis. There may be a transient increase in white cell count. CSF pressure may be increased, the protein content of CSF is usually increased. Pleocytosis is present. The best way to fix the diagnosis is to isolate the virus from blood. But, this is very uncommon in this disease. The serological study might prove the diagnosis. The accuracy of serological study is decreased by the prior presence of other related viruses. Early Ig M antibody reactive in neutralisation and haemagglutination-inhibition test only is a feature. Ig G antibody may be detected after 8-10 days which is reactive in complement fixation test.

TREATMENT AND PROGNOSIS :

The treatment is intensive supportive care. There is no specific treatment. Artificial control of hyperthermia should be attempted. Antipyretics are hardly of any use. So, cold

sponging should be done. Adequate nutrition should be given by intravenous route. Steroids have to be used in most of these deteriorating cases. Antibiotics are used in case of secondary infection. The mortality rate in the adult is about 80%. It is about 30% in the children, but permanent sequelae like UMN paralysis, cerebellar syndromes, personality changes and mental deterioration are more common in them.

PREVENTION :

It is now clear that prevention is the only way out. The following preventive methods are attempted. The first one is "Live attenuated vaccine". It was used with success in pigs in Japan. But this is too dangerous to be used in human for fear of overt infection from the vaccine itself. The second one is "Formalised vaccine". Some success has been reported with this in Japan. But the poor people of this area has yet to get it. The third one is "Passively acquired antibody". Due to sophisticated method and the cost involved, it is out of reach of the people of this part of the world. The one and only way of prevention that is feasible here is specific vector control. His Majesty's Govt. has gone all out towards this end.

— o —

Now On the Scene But for all types of tough

PHENCODIN

COUGH SYRUP

ROYAL DRUGS LTD.
NEWARK.

RD-AMPI

*Published by the Nepal Medical Association, Siddhi Sadan, Exhibition Road
Printed at Naveen Press, Dugambahil, (New Road) Kathmandu, Nepal.*