Evidence Of Leishmaniasis In Nepal

Dr. D. D. Joshi*

Introduction:

Leishmaniasis is a parasitic zoonosis. Almost all forms of leishmaniasis are zoonosis. There are two types of leishmaniasis – one cutaneous leishmaniasis and other visceral leishmaniasis. Both types are caused by a flagellate protozoan of the genus Leishmania which occur in vertebrate reservoir animals, including man, in the aflagellate form (leishmanial) of amastigote form, while in the vector insects Phlebotomus spp. it has the flagellate (leptomonad) or promastigote form.

Visceral leishmaniasis, which is also called kala-azar or dum-dum fever or infantile splenic fever, is caused by Leishmania donovani which is a viscerotropic flagellated protozoan.

Cutaneous leishmaniasis of oriental sore, or Aleppo boil is caused i.e. L. mexicana mexicana, and L. Grasilensis brasiliensis. L. Peruviana. L. tropica major, and L. tropica minor (Fifth Annual on TDR 1981).

Disease Distribution in man and Animals:

Cutaneous and visceral leishmaniasis in humans and animals occur in the Americas from southern Mexico to northern, Argentina, Asia the Middle East, the Mediterranean coast and Africa. On a global basis this is prevalent but very poorly reported and generally underestimated, (Acha et al 1980).

In Nepal this disease is prevalent both in human and animals but it is very poorly

* Chief Zoonotic Diseases Control Section, Epidemiology & Statistics Division, Department of Health Services and Member-Secretary, Nepal Medical Research Committee.

reported and underestimated. Stray cases of Kala-azar are occurring in Siraha, Dhanusha, Mahotari, Banke and Kailali districts of Nepal. There is one case diagnosed clinically as well as laboratorically in Kanti Hospital, Kathmandu (Shrestha 1982). This calls for active epidemiological surveillance to prevent this disease situation in Nepal. Sandflies (*Phlebotomus* spp.) which transmit this disease from animal to man are also present in the endemic districts of Nepal.

In India both types of leishmaniasis is present in Asam, Bihar, U.P., Rajasthan, West-Bangal and areas of river Ganga. In Bihar this disease is existing since 1933. From 1970 onwards cases of Kala-azar started reappearing in Bihar and in 1977 the estimated cases reached about 100,000 with approximately 4500 deaths. From 1980-82 this disease appeared more severely in many districts of Bihar which are adjacent to Siraha, Dhanusha Mahotari & Saptari districts (Sanyal et al 1979). The investigations revealed the first presence of a zoonotic focus of cutaneous leishmaniasis in Bikaner district of Rajasthan in 1973 (Sharma et al 1973 and Mohan Krishna et al 1973).

Epidemiological Transmission Cycle:

The reservoirs of visceral leishmaniasis (Kala-azar) are dogs and other canine species and to a lesser extent man himself. The infection phlebotomine insects (sandflies). Then the infected sandfly bites man and transmits the disease. Occasionally transmission is from man to man by the bites of sandfly. (see Figure 1)

The reservoirs of cutaneous leishmaniasis are wild animals particularly rodents. The infection is transmitted from one animal to another by phlebotomine flies of the genus Lutzomyia. Man is infected accidentally by the bites of these flies when he enters enzootic areas in the forest (See Figure 2).

In Dhanusha district, Nepal Malaria Eradication office has collected sandflies during the year 1982 when there was an out-break of Kala - azar in the area. But no detail study has been carried out. Sandflies (phlebotomus argentipes) are important vector of Kala-azar in Bihar, India which is bordered with Dhanusha district of Nepal.

Reported Cases of Kala-azar In Nepal:

Cases of Kala-azar started being reported and admitted from 1981 onwards from

Janakpur Zonal Hospital, Dhanusha. In that year 1981 there were 5 cases clinically confirmed and treated in the hospital, and in 1982 there were 32 cases reported and 7 hospitalized (See Table I). Besides many patients went to India for treatment. Clinical cases of Kala-azar were also reported from Siraha, Saptari, Mahotari, Banke, Kailali and Kanti Hospital in different years.

Zoonotic Diseases Control Section, Epidemiology and Statistics Division of the Department of Health Services was asked to do some epidemiological survey of this disease in that area. Then the project proposed has been prepared and approved by Nepal Medical Research Committee, Ministry of Health for implementation. The project has been sent to TDR Division WHO/H.Q. Geneva. Preliminary epidemiological survey has been already started now by this section.

Acknowledgement:

The author would like to thank Dr. Pitambar Jha, then Civil Surgeon of Janakpur Zonal Hospital for his kind cooperation and help.

Figure 1

Epidemiological Transmission Cycle of visceral Leishmaniasis (Kala azar).

Figure 2

Epidemiological Transmission Cycle of Cutaneous Leishmaniasis

Table 1

Clinical Cases of Kala-azar Admitted in Janakpur

Zonal Hoapital, Dhanusha (1981-1982)

Cases No.	Age	Sex	Address	Cured	Death
01.	12	F	Kapleswar	Cured	-----
02.	8	M	Kamalakhoch	Cured	-----
03.	30	M	Janakpur	Cured	-----
04.	22	M	Kapleswar	Left hospital without medical advise(LAMA)	-----
05.	45	M	Janakpur	Cured	-----
06.	16	F	Bhimban	-----	Died
07.	5	M	Basantpur	Cured	-----
08.	8	F	Kapleswar	LAMA	-----
09.	24	F	Hardiga	Cured	-----
10.	22	F	Bherdar	Cured	-----
11.	10	M	Lohna	Cured	-----
12.	3	F	Bakey	Cured	-----

REFERENCES

1. Acha, P.N and Szyfres, B. (1980). Zoonoses and Communicable Diseases Common to Man and Animals. Visceral Leishmaniasis. PP: 413-418. Pan American Health Organization/WHO, Washington D.C. U S A.
2. Fifth Annual Report (1981). UNDP / World Bank / WHO. Special Programme for Research and Training in Tropical Diseases.
3. Mohan Krishan, Suri, J.C. and Madan Meera (1973). Studies on Cutaneous Leishmaniasis in India. II Behaviour of Leishmania Tropica strain isolated from dogs in various experimental animals. J. Com. Dis. 5(3): 143 - 148.

4. Sanyal, R.K., Alam, S.M. and Wottal, B.L. (1979). *Some observations on Epidemiology of current out break of Kala-azar in Bihar.* J. com Dis. 11(4): 170 - 184.
5. Sanyal, R.K., Banerjee, D.P., Ghosh, T. K., Ghosh, J. N., Misra, B. S., Roy, Y. P. and Rao, C. K. (1979). *A Longitudinal Review of Kala-azar in Bihar.* J. Com. Dis. 11(4) 149 - 169.
6. Sharma, M.I. D., Suri, J. C. and Lalra, N. L. (1973). *Studies on Cutaneous Leishmaniasis I, A note on the current status of III. Detection of a Zoonotic focus of cutaneous Leishmaniasis in Rajasthan.* J. Com. Dis. 5(3) : 149 - 153.
8. Shrestha, Y. B. (1982) . *Kala-azar in Kanti Hospital Nepal J. Vol. I, No. 1 , PP: 212-216.*